

Lecture 13: 02/22/2022

Lecturer: Xiaorui Sun

Scribe: Sriman
Cherukuru

1 Last Lecture's Review

Last Lecture: Graph decomposition

- Utilization of nice properties on good edges (intra-component)
- Upper bound on the number of bad edges (inter-component)

This Lecture: Graph sparsification (Edge, Vertex sparsification)

- Preserve important properties of the original graph
- Bounding the size of sparsified graph for computation efficiency

2 Graph sparsification

2.1 Edge sparsification

$$G = (V, E) \rightarrow H = (V, E')$$

Such that: property on $H \approx$ property on G , and $|E'| \ll |E|$. Eg.,
Spanning forest for graph connectivity property.

2.2 Vertex sparsification

$$G = (V, E), (\text{terminal set}) T \subset V \rightarrow H = (V', E')$$

Such that: $T \subset V'$, property on T in $H \approx$ property on T in G , $|V'| \ll |V|$, $|E'| \ll |E|$. Eg.,
weighted graph for distance property.

3 VERTEX SPARSIFICATION FOR MIN-CUT PROPERTY



3 Vertex sparsification for min-cut property

3.1 Problem

Given $G = (V, E)$, $T \subset V$, construct a vertex sparsifier $H = (V', E')$ such that: $T \subset V'$, and for all partitions of T into $(T', T \setminus T')$:

$$\text{mincut}_H(T', T \setminus T') = \text{mincut}_G(T', T \setminus T').$$

With additional parameter of threshold c :

If $\text{mincut}_G(T', T \setminus T') \leq c$, then $\text{mincut}_H(T', T \setminus T') = \text{mincut}_G(T', T \setminus T')$.

If $\text{mincut}_G(T', T \setminus T') > c$, then $\text{mincut}_H(T', T \setminus T') > c$.

3.2 The algorithm

Definition 1 (Cut containment set) A set of edges E' in G is called a cut containment set if $\forall (T', T \setminus T')$ with $\text{mincut}_G(T', T \setminus T') \leq c, \exists$ such a min-cut with edges C and $C \subset E'$.

Definition 2 (Intersecting-all-mincut set (old definition)) Given $G = (V, E)$, terminal set $T \subset V$ and threshold c , we say a set of edges F in G is a set of edges intersecting all min terminal cuts if $\forall (T', T \setminus T')$ with $\text{mincut}_G(T', T \setminus T') \leq c, \exists$ such a min-cut with edges C and $C \cap F \neq \emptyset$.

Definition 3 (Intersecting-all-mincut set (new definition)) Given $G = (V, E)$, terminal set $T \subset V$ and threshold c , we say a set of edges F in G is a set of edges intersecting all min terminal cuts if after removing F from G , $\forall (T', T \setminus T')$ with $\text{mincut}_G(T', T \setminus T') \leq c$, no connected component in $G \setminus F$ contains all the edges of a terminal min-cut.

Theorem 4 For vertex sparsification with respect to the min-cut property with threshold parameter c , there exists a sparsifier of size $|T| \cdot c^{O(c)}$.


(Can be improved to $|T| \cdot c^3 \cdot \log^4 n$.)

Suppose we have a cut containment set E' , we can construct a vertex sparsifier H from E' in the following way:

- Remove edges in E' from G , breaking G into several connected components.
- Shrink each connected component into 1 super-vertex in H .

- Connect vertices in H with corresponding edges in E' .
- If terminal vertices u, v are merged into the same connected component (thus the terminal nodes are lost in H), add u, v as vertices in H , connect each with $c+1$ parallel edges to the super-vertex corresponding to the connected component they're in. (There were no min-cut of size $\leq c$ between them in G .)

Resulting H has number of edges (and vertices) $\leq |E'| + |T| \cdot (c+1)$. Next, we need to upper-bound size of E' .



Claim 5 Given $G, T, c, \exists E'$ st. $|E'| = |T| \cdot c^{O(c)}$.

Proof idea: recursively build E' using intersecting-all-set F :

- Initialize $E' = \emptyset$.
- Find intersecting-all-set F_c in G for given $T = T_0$ and c .
- Remove all edges in F_c from G , add them to E' , and add endpoints of edges in F_c to terminals:
 $G \leftarrow G \setminus F_c, E' \leftarrow E' \cup F_c, T_1 \leftarrow T_0 \cup \{u, v \mid (u, v) \in F_c\}$.
- Continue to find intersecting-all-set F_{c-1} with updated G and T_1 , until $c = 1$.
- Return E' (which is $\bigcup_{i=1}^c F_c$).

Claim 6 Given G, T, c , the intersecting-all-set F_c is of size $|F_c| \leq |T| \cdot c$. (Each terminal vertex is incident to at most c edges that are on any qualifying min terminal cut.)

Proof of Claim 5 by induction:

Base case: $|F_c| \leq |T| \cdot c$.

Step 1: $|T_1| \leq |T| + 2|F_c| \leq (1+2c) \cdot |T| = O(c) \cdot |T|$, and $|F_{c-1}| \leq |T_1| \cdot c = O(c^2) \cdot |T|$.

General induction step: $|T_i| \leq |T_{i-1}| + 2|F_{c-i+1}| = O(c^i) \cdot |T|$, $|F_{c-i}| = O(c^{i+1}) \cdot |T|$. Finally, $|F_1| = O(c^c) \cdot |T|$.

$|E'| = |\bigcup_{i=1}^c F_c| = c^{(c)} \cdot |T|_0$.